



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Vertical transmission of *Mycoplasma pneumoniae* infection

Huber, Benedikt M ; Meyer Sauter, Patrick M ; Unger, Wendy W J ; Hasters, Paul ; Eugster, Marcel R ; Brandt, Simone ; Bloemberg, Guido V ; Natalucci, Giancarlo ; Berger, Christoph

Abstract: *Mycoplasma pneumoniae* is a significant cause of pneumonia in school-aged children and young adults. We report a case of neonatal *M. pneumoniae* pneumonia in a preterm child manifesting in the first hours of life. Vertical transmission was demonstrated by the detection of *M. pneumoniae* in inflamed placental tissue indicating chorioamnionitis.

DOI: <https://doi.org/10.1159/000490610>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-153241>

Journal Article

Accepted Version

Originally published at:

Huber, Benedikt M; Meyer Sauter, Patrick M; Unger, Wendy W J; Hasters, Paul; Eugster, Marcel R; Brandt, Simone; Bloemberg, Guido V; Natalucci, Giancarlo; Berger, Christoph (2018). Vertical transmission of *Mycoplasma pneumoniae* infection. *Neonatology*, 114(4):332-336.

DOI: <https://doi.org/10.1159/000490610>



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
Main Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Vertical transmission of *Mycoplasma pneumoniae* Infection

Huber, Benedikt M; Meyer Sauter, Patrick M; Unger, Wendy WJ; Hasters, Paul; Eugster, Marcel R;
Brandt, Simone; Bloemberg, Guido V; Natalucci, Giancarlo; Berger, Christoph

DOI: <https://doi.org/10.1159/000490610>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-168060>

Journal Article

Originally published at:

Huber, Benedikt M; Meyer Sauter, Patrick M; Unger, Wendy WJ; Hasters, Paul; Eugster, Marcel R; Brandt, Simone; Bloemberg, Guido V; Natalucci, Giancarlo; Berger, Christoph (2018). Vertical transmission of *Mycoplasma pneumoniae* Infection. *Neonatology*, 114(4):332-336.

DOI: <https://doi.org/10.1159/000490610>

Vertical Transmission of *Mycoplasma pneumoniae* Infection

Benedikt M. Huber^a Patrick M. Meyer Sauter^b Wendy W.J. Unger^c Paul Hasters^a
Marcel R. Eugster^d Simone Brandt^e Guido V. Bloemberg^f Giancarlo Natalucci^a
Christoph Berger^{b,f}

^aDepartment of Neonatology, University Hospital Zurich, Zurich, Switzerland; ^bDivision of Infectious Diseases and Hospital Epidemiology, and Children's Research Center (CRC), University Children's Hospital Zurich, Zurich, Switzerland;

^cLaboratory of Pediatrics, Division of Pediatric Infectious Diseases and Immunology, Erasmus MC University Medical Center – Sophia Children's Hospital, Rotterdam, The Netherlands; ^dUnilabs Duebendorf, Molecular Diagnostics, Duebendorf, Switzerland;

^eInstitute of Pathology and Molecular Pathology, University Hospital Zurich, Zurich, Switzerland; ^fInstitute of Medical Microbiology, University of Zurich, Zurich, Switzerland

Established Facts

- *Mycoplasma pneumoniae* causes pneumonia predominantly in school-aged children and young adults.
- Neonatal pneumonia associated with *M. pneumoniae* has been very rarely reported.

Novel Insights

- Vertical transmission of *Mycoplasma pneumoniae* infection was demonstrated in this case, the first time with the detection of *M. pneumoniae* by PCR and immunohistochemistry in placental tissue.
- *M. pneumoniae* can be considered as possible cause of congenital pneumonia in addition to other mycoplasmas (*M. hominis*) and ureaplasmas (*U. urealyticum* and *U. parvum*).

Keywords

Congenital infection · *Mycoplasma pneumoniae* · Neonatal pneumonia · Vertical transmission

Abstract

Mycoplasma pneumoniae is a significant cause of pneumonia in school-aged children and young adults. We report a case of neonatal *M. pneumoniae* pneumonia in a preterm child manifesting in the first hours of life. Vertical transmission was demonstrated by the detection of *M. pneumoniae* in inflamed placental tissue indicating chorioamnionitis.

© 2018 S. Karger AG, Basel

Introduction

Mycoplasma pneumoniae colonizes the upper respiratory tract [1] and causes pneumonia predominantly in school-aged children and young adults [2]. In contrast, other mycoplasmas and ureaplasmas colonize the urogenital tract, among which *M. hominis*, *U. urealyticum*, and *U. parvum* may cause ascending intrauterine infection that can lead to adverse pregnancy outcomes and/or

B.M.H. and P.M.M.S. contributed equally to this work.

neonatal pneumonia [3, 4]. Here, we present a preterm infant with severe neonatal *M. pneumoniae* pneumonia acquired by vertical infection.

Case Report

A preterm male neonate weighing 1,500 g was delivered by a 30-year-old woman at 29 4/7 weeks of gestation by cesarean section because of recurrent vaginal bleedings and premature contractions for 3 days. Antenatal steroid administration had been completed. The Apgar score was 2, 4, and 9 at 1, 5, and 10 min, respectively, and the umbilical artery pH was 7.31. He developed a severe respiratory distress syndrome (RDS) in the first hour of life requiring mechanical ventilation and surfactant administration. Chest X-ray showed a granular appearance of both lungs with air bronchograms (Fig. 1a). Empiric antibiotic treatment with amoxicillin and gentamicin was immediately started. After 24 h, extubation was achieved and followed by nasal continuous positive airway pressure treatment. Antibiotic treatment was discontinued based on negative blood cultures and C-reactive protein within the normal range. Secondary respiratory distress developed on the second day of life (DOL) and necessitated re-intubation until DOL 4 and again from DOL 6 to 11. Chest X-ray on DOL 6 revealed multifocal opacifications and consolidations (Fig. 1b). Because of the atypical RDS presentation, an extensive diagnostic workup was performed: cultures from blood and tracheal aspirate were repeatedly negative for bacteria, as well as for fungi, as were cultures from urine for cytomegalovirus. There were no signs and symptoms of multiorgan involvement. Blood cell count showed a leukocytosis of $41 \times 10^9/\text{L}$ after birth, which increased to a maximum of $97 \times 10^9/\text{L}$ on DOL 2 and consisted of mainly neutrophils, including immature granulocytes. There was no evidence of leukemia or transient myeloproliferative disorder. C-reactive protein remained normal over the course of disease.

The unclear situation led to a detailed review of the medical history during pregnancy: the mother recalled a mild respiratory tract infection with intractable cough at 20 gestational weeks lasting for a week, but this was left untreated. The diagnostic workup in the neonatal tracheal aspirate was extended by *M. pneumoniae*-specific PCR as previously described [5]: *M. pneumoniae* DNA could be detected in tracheal aspirate on DOL 3 and in a second sample from nasopharyngeal aspirate after extubation on DOL 4. No DNA of *M. hominis*, *M. genitalium*, or *Ureaplasma* spp. was found in the tracheal aspirate by PCR, performed as described previously [6, 7]. Treatment with erythromycin was initiated orally on DOL 4 (50 mg/kg/dose 4 times a day) and switched to intravenous application from DOL 7 to DOL 18 (40 mg/kg/dose 4 times a day). Erythromycin treatment was paralleled by a steady and sustainable improvement of clinical and radiographic findings. Chest X-ray on DOL 9 returned almost to normal. On DOL 22, the white blood cell count was normal and serological testing using an enzyme-linked immunosorbent assay (Serion GmbH, Würzburg, Germany) revealed *M. pneumoniae*-specific immunoglobulin (Ig) M and IgG antibodies of <5 U/mL (cutoff 17 U/mL) and 65 U/mL (cutoff 15 U/mL), respectively. Nasal continuous positive airway pressure treatment was followed until DOL 23 and supplemental oxygen administered until DOL 30, defining mild bronchopulmonary dysplasia. The infant was discharged with 8 weeks of age at 37 6/7 weeks postmenstrual age. A 1-month follow-up was uneventful.

Maternal serum obtained 2 weeks after birth was tested positive for *M. pneumoniae*-specific IgM (93 U/mL; cutoff 17 U/mL) and IgG (>200 U/mL; cutoff 30 U/mL), indicating a recent infection. Prepartal maternal swabs from the cervix uteri were negative by PCR for DNA of *Chlamydia trachomatis* and *Neisseria gonorrhoeae*, and also vaginal swab cultures were negative. Histological examination of the placenta showed distinct chorioamnionitis and vasculitis with infiltration of neutrophils into the chorioamniotic mesoderm layer and amnion (Fig. 1c, e). Placental tissues embedded in paraffin were tested positive for *M. pneumoniae* DNA by PCR and *M. pneumoniae* antigens by immunohistochemistry (Fig. 1d, f). Placental tissue was tested negative for DNA of *Ureaplasma* spp., *M. hominis*, and *M. genitalium* by PCR. Control placental tissues without chorioamnionitis and chorioamnionitis of other origin were tested negative for *M. pneumoniae* antigens by immunohistochemistry (data not shown).

Discussion

Congenital pneumonia arises from direct mucosal seeding from infected amniotic fluid (chorioamnionitis), which is caused by hematogenous transplacental infection or ascending infection across the chorioamniotic membranes [8]. Maternal vaginal colonization is a key risk factor for an ascending intrauterine infection and/or perinatal infection during passage through the birth canal [8]. Mycoplasmas are primarily mucosal pathogens, among which genital mycoplasmas and ureaplasmas colonize the urogenital tract. *M. hominis*, *U. urealyticum*, and *U. parvum* are also associated with neonatal pneumonia [3, 4]. In contrast, *M. pneumoniae* is known to exclusively colonize the respiratory tract [3]. The question arises whether chorioamnionitis in our case was caused by a so far not reported ascending infection or rather by spread from respiratory tract infection through the bloodstream to the placenta. In line with the latter, *M. pneumoniae* has been reported to disseminate in the bloodstream during or after a respiratory tract infection and to cause extrapulmonary manifestations [9–12]. The mother indeed experienced a cough around 8 weeks before birth and the serology after birth confirmed a recent *M. pneumoniae* infection. This respiratory infection may have led to invasive infection and spread of *M. pneumoniae* to the placenta. In fact, infections with *M. hominis*, *U. urealyticum*, or *U. parvum* as potential cause of adverse pregnancy outcome and/or neonatal pneumonia were excluded. Thus, the diagnosis of vertical *M. pneumoniae* infection in our case is established as follows: (1) maternal respiratory tract infection at 20 gestational weeks with strongly positive *M. pneumoniae* serology 2 weeks after birth; (2) detection of *M. pneumoniae* in placental tissue by PCR and immuno-




Fig. 1. **a** Chest X-ray, on day of life (DOL) 1, showing a diffuse interstitial pattern with granular appearance and air bronchograms of both lungs. **b** Chest X-ray, on DOL 6, showing reticular (bullous) multifocal opacifications and consolidations. **c, e** Chorioamnionitis with infiltration of neutrophils (arrows) into the chorioamniotic mesoderm layer and amnion. Hematoxylin and eosin stain. Original magnification $\times 100$ (**c**) and $\times 400$ (**e**). **d, f** Immunohistochemical analysis of placental tissue performed by using a biotinylated polyclonal anti-*M. pneumoniae* antibody (Thermo Scientific, Waltham, MA, USA) and an avidin-biotin-peroxidase complex with 3,3'-diaminobenzidine tetrahydrochloride chromogenic substrate showing positive staining in the chorioamniotic mesoderm layer and amnion. Original magnification $\times 100$ (**d**) and $\times 400$ (**f**).

Table 1. Overview of published cases on *M. pneumoniae* infections in neonates including the case report

Case No.	Ref.	GA	BW, g	Signs/symptoms	Onset	Chest X-ray	<i>M. pneumoniae</i> PCR	<i>M. pneumoniae</i> serology	Placenta	Pregnancy	Maternal <i>M. pneumoniae</i> serology	Treatment	Outcome
1	10	38	3,550	Respiratory failure, mucus	DOL 1	Pneumonia, pneumo-thorax	+ (TA)	-	NA	Mother: URTI at GA 32–38 weeks Sister: <i>M. pneumoniae</i> PCR+ at GA 32 weeks	NA	Amoxicillin and netilmicin IV	Normal
2	11	30	1,685	Respiratory failure, severe BPD	DOL 1	Pneumonia	+ (NPA)	-	Chorioamnionitis and cord vasculitis; <i>M. pneumoniae</i> PCR+	Mother: URTI at GA 25 weeks	+ (seroconversion during pregnancy)	Erythromycin PO 7 days, azithromycin PO 28 days	Demise at PMA 44 weeks; severe BPD
3	12	39	NA	Fever, crying, respiratory distress, feeding problems	DOL 14	Pneumonia	-	+ (IgM and IgG, seroconversion)	NA	NA	-	Erythromycin PO 14 days	Normal
4	Case report	29	1,500	Respiratory failure	DOL 1	Pneumonia	+ (TA + NPA)	+ (IgG)	Chorioamnionitis and cord vasculitis; <i>M. pneumoniae</i> PCR+	Mother: URTI at GA 20 weeks	+ (IgM and IgG postpartal)	Erythromycin PO/IV 14 days	Mild BPD

BW, birth weight; BPD, bronchopulmonary dysplasia; DOL, day of life; GA, gestational age (in completed weeks); *M. pneumoniae*, *Mycoplasma pneumoniae* IV, intravenous; NPA, nasopharyngeal aspirate; NA, not available; PCR, polymerase chain reaction; PMA, postmenstrual age; PO, orally; TA, tracheal aspirate; URTI, upper respiratory tract infection.

histochemistry; (3) detection of *M. pneumoniae* by PCR from neonatal respiratory specimens on DOL 3 and 4; and (4) neonatal pneumonia manifesting in the first hours of life and presenting as atypical RDS.

Invasive *M. pneumoniae* infection is rare [10], and vertical transmission of *M. pneumoniae* infection has been very rarely reported. To our knowledge, 3 cases of neonatal pneumonia associated with *M. pneumoniae* have been published so far (Table 1) [13–15]. Vertical transmission of *M. pneumoniae* infection has been suggested in 2 pre-term neonates with either rapidly or slowly progressing respiratory failure requiring mechanical ventilation immediately after birth. A vertical route of transmission was confirmed in 1 case with the detection of *M. pneumoniae* DNA by PCR in placental tissue. We additionally showed by immunohistochemical analysis that *M. pneumoniae* is present in the placenta.

Interestingly, anti-*M. pneumoniae* IgG, but not IgM, was detected in the neonate on DOL 22. The detection of specific IgG is complicated by transplacental transfer of maternal antibodies. In contrast, the detection of IgM is very specific to the fetal compartment because IgM does not cross the placenta. However, the neonate's immune system may not mount an antibody response as effective as adults [16], and, most importantly, a negative IgM result does not exclude congenital infection [17, 18]. Further, the antibody response to *M. pneumoniae* is complex [19]. We present only the second case of neonatal pneumonia associated with *M. pneumoniae*, in which the antibody response was assessed (Table 1). One might speculate that the presence or absence of *M. pneumoniae*-specific IgM may discriminate between perinatal infection after birth (case 3, Table 1) and congenital infection (present case).

This case demonstrates that *M. pneumoniae* can be considered as possible cause of congenital pneumonia in addition to other “atypical” organisms. The route of transmission of *M. pneumoniae* is vertical infection after dissemination of the bacteria following maternal respiratory tract infection. Chorioamnionitis likely induced premature birth, but it remains unclear whether *M. pneumoniae* triggered also bronchopulmonary dysplasia as reported for ureaplasmas.

Acknowledgments

We thank the technicians of the Institute of Medical Microbiology, University of Zurich, for their assistance with microbiological analyses, and Silvia C. Estevão (Laboratory of Pediatrics, Erasmus MC – Sophia Children's Hospital, Rotterdam) for her assistance

with immunohistochemical analyses. P.M.M.S. was supported by grants of the Promedica and Starr International Foundation, and a Fellowship Award of the European Society for Paediatric Infectious Diseases (ESPID), all outside the submitted work.

Statement of Ethics

Informed consent has been obtained.

Disclosure Statement

There is nothing to disclose.

Funding Source

There is no funding to disclose.

Author Contributions

B.M.H., P.H., G.N.: patient care and routine diagnostic work-up; P.M.M.S., W.W.J.U., M.R.E., S.B., G.V.B., C.B.: microbiological, histological, immunohistochemical analyses; P.M.M.S., C.B.: writing the manuscript; C.B.: coordinating the work; all authors: critically reviewing the manuscript.

References

- Spuesens EB, Fraaij PL, Visser EG, Hoogenboezem T, Hop WC, van Adrichem LN, Weber F, Moll HA, Broekman B, Berger MY, van Rijsoort-Vos T, van Belkum A, Schutten M, Pas SD, Osterhaus AD, Hartwig NG, Vink C, van Rossum AM: Carriage of *Mycoplasma pneumoniae* in the upper respiratory tract of symptomatic and asymptomatic children: an observational study. *PLoS Med* 2013;10:e1001444.
- Jain S, Williams DJ, Arnold SR, Ampofo K, Bramley AM, Reed C, Stockmann C, Anderson EJ, Grijalva CG, Self WH, Zhu Y, Patel A, Hymas W, Chappell JD, Kaufman RA, Kan JH, Dansie D, Lenny N, Hillyard DR, Haynes LM, Levine M, Lindstrom S, Winchell JM, Katz JM, Erdman D, Schneider E, Hicks LA, Wunderink RG, Edwards KM, Pavia AT, McCullers JA, Finelli L; CDC EPIC Study Team: Community-acquired pneumonia requiring hospitalization among US children. *N Engl J Med* 2015;372:835–845.
- Waites KB, Katz B, Schelonka RL: Mycoplasmas and ureaplasmas as neonatal pathogens. *Clin Microbiol Rev* 2005;18:757–789.
- Viscardi RM: *Ureaplasma* species: role in neonatal morbidities and outcomes. *Arch Dis Child Fetal Neonatal Ed* 2014;99:F87–F92.
- Hardegger D, Nadal D, Bossart W, Altwegg M, Dutly F: Rapid detection of *Mycoplasma pneumoniae* in clinical samples by real-time PCR. *J Microbiol Methods* 2000;41:45–51.
- Pascual A, Jaton K, Ninet B, Bille J, Greub G: New diagnostic real-time PCR for specific detection of *Mycoplasma hominis* DNA. *Int J Microbiol* 2010;2010:317512.
- Xiao L, Glass JI, Paralanov V, Yooseph S, Cassell GH, Duffy LB, Waites KB: Detection and characterization of human *Ureaplasma* species and serovars by real-time PCR. *J Clin Microbiol* 2010;48:2715–2723.
- Hooven TA, Polin RA: Pneumonia. *Semin Fetal Neonatal Med* 2017;22:206–213.
- Neimark H, Gesner M: Is *Mycoplasma pneumoniae* adherence to erythrocytes a factor in extrapulmonary dissemination? *PLoS Pathog* 2010;6:e1001219.
- Meyer Sauter PM, Jacobs BC, Spuesens EB, Jacobs E, Nadal D, Vink C, van Rossum AM: Antibody responses to *Mycoplasma pneumoniae*: role in pathogenesis and diagnosis of encephalitis? *PLoS Pathog* 2014;10:e1003983.
- Narita M: Pathogenesis of neurologic manifestations of *Mycoplasma pneumoniae* infection. *Pediatr Neurol* 2009;41:159–166.
- Narita M: Pathogenesis of extrapulmonary manifestations of *Mycoplasma pneumoniae* infection with special reference to pneumonia. *J Infect Chemother* 2010;16:162–169.
- Ursi D, Ursi JP, Ieven M, Docx M, Van Rempts P, Pattyn SR: Congenital pneumonia due to *Mycoplasma pneumoniae*. *Arch Dis Child Fetal Neonatal Ed* 1995;72:F118–F120.
- Srinivasjois RM, Kohan R, Keil AD, Smith NM: Congenital *Mycoplasma pneumoniae* pneumonia in a neonate. *Pediatr Infect Dis J* 2008;27:474–475.
- Kumar S, Maria A, Saigal SR, Maheshwari M: *Mycoplasma pneumoniae* as a cause of non-resolving pneumonia in a neonate. *J Med Microbiol* 2010;59:731–732.
- Hassan J, Dooley S, Hall W: Immunological response to cytomegalovirus in congenitally infected neonates. *Clin Exp Immunol* 2007;147:465–471.
- Herremans T, Kortbeek L, Notermans DW: A review of diagnostic tests for congenital syphilis in newborns. *Eur J Clin Microbiol Infect Dis* 2010;29:495–501.
- Coll O, Benoist G, Ville Y, Weisman LE, Botet F, Anceschi MM, Greenough A, Gibbs RS, Carbonell-Estrany X; WAPM Perinatal Infections Working Group: Guidelines on CMV congenital infection. *J Perinat Med* 2009;37:433–445.
- Meyer Sauter PM, de Bruijn ACJM, Graça C, Tio-Gillen AP, Estevão SC, Hoogenboezem T, Hendriks RW, Berger C, Jacobs BC, van Rossum AMC, Huizinga R, Unger WWJ: Antibodies to protein but not glycolipid structures are important for host defense against *Mycoplasma pneumoniae*. Submitted.